

公募研究「初期特異点のない新しい宇宙モデルとその観測的検証」

Generic instabilities of non-singular cosmologies in second-order theories: **A no-go theorem**



Tsutomu Kobayashi
Rikkyo University

小林努
立教大学

Based on

TK, PRD94 (2016) 04351 [arXiv:1606.05831]


Shingo Akama & TK, PRD accepted,

[arXiv:1701.02926] w/ 赤間進吾(立教大学)

Introduction

- **Inflation** is an almost perfect scenario, but
 - Initial singularity Borde & Vilenkin gr-qc/9612036
 - Trans-Planckian problem Martin & Brandenberger hep-th/0005209
- **Non-singular cosmologies**: bounce, Galilean Genesis, ...
 - ~~Null energy condition (NEC)~~
$$\dot{H} = -4\pi G (\rho + p) > 0$$
 - Stability?

Battefeld & Peter 1406.2790
Brandenberger & Peter 1603.05834



NEC and stability

- Quadratic Lagrangian for curvature perturbation

$$\mathcal{L} = a^3 \left[\mathcal{G}_S(t) \dot{\zeta}^2 - a^{-2} \mathcal{F}_S(t) (\vec{\nabla} \zeta)^2 \right]$$

Ghost/gradient instabilities if $\mathcal{G}_S < 0$ / $\mathcal{F}_S < 0$



- $P(\phi, X)$ cosmologies are unstable if NEC is violated:

$$X := -\frac{1}{2}(\partial\phi)^2 \quad \nearrow \quad \mathcal{F}_S = (8\pi G)^{-1}(-\dot{H})/H^2$$

- Galileon-type 2nd-order theories,

$$\mathcal{L} = \frac{R}{2\kappa} + G_2(\phi, X) - G_3(\phi, X)\square\phi + \dots$$

admit stable NEC-violating solutions, because $\mathcal{F}_S$ and $\mathcal{G}_S$ are not correlated with the sign of $\dot{H}$

Stable non-singular cosmologies?


- Stable NEC-violating phases are possible
- But, known examples exhibit **instability at some moment in the entire history**
- Typically,
 - gradient instability at the bounce point
 - at the transition from NEC-violating phase to another
 - or at some moment after the bounce



Creminelli, *et al.* 1007.0027; Qui, *et al.* 1108.0593; Easson, *et al.* 1109.1047; Osipov & Rubakov 1303.1221; Cai, *et al.* 1206.2382; Koehn, *et al.* 1310.7577; Pirtskhalava, *et al.* 1410.0882; TK, Yamaguchi, Yokoyama 1504.05710;

Example 1 (Bounce)

Koehn, Lehnert, Ovrut 1310.7577




Gradient instability at the bounce

Example 2 (Galilean Genesis)

Pirtskhalava, *et al.* 1410.0882


TK, Yamaguchi, Yokoyama 1504.05710



Gradient instability at the transition from Galilean Genesis to inflation

Example 3 (Bounce)

Easson, Sawicki & Vikman 1109.1047



Stable bounce, but eventually $c_s^2 < 0$

Question to be addressed

- Is the appearance of gradient instabilities generic or model-dependent nature?

- A partial answer was given by Libanov, Mironov & Rubakov 1605.05992

- All non-singular cosmological solutions in

$$\mathcal{L} = \frac{R}{2\kappa} + G_2(\phi, X) - G_3(\phi, X)\Box\phi$$

are plagued with gradient instabilities

- This talk: *The no-go theorem can be extended to the most general 2nd-order scalar-tensor theory (Horndeski) and to a wide class of theories with multiple scalar fields*

Horndeski theory

$$\mathcal{L} = G_2(\phi, X) - G_3(\phi, X)\Box\phi + G_4(\phi, X)R \\ + G_{4,X} [(\Box\phi)^2 - (\nabla_\mu\nabla_\nu\phi)^2] + G_5(\phi, X)(\cdots)$$

Horndeski (1974); Deffayet, *et al.* 1103.3260;
TK, Yamaguchi, Yokoyama 1105.5723

- The most general scalar-tensor theory with 2nd-order field equations
- Einstein + canonical scalar, Brans-Dicke, $f(R)$, k-essence, covariant Galileons, dilatonic Gauss-Bonnet, ... as specific cases

Perturbations in Horndeski

TK, Yamaguchi, Yokoyama 1105.5723

■ Tensor perturbations

$$\mathcal{L} = \frac{a^3}{8} \left[\mathcal{G}_T(t) \dot{h}_{ij}^2 - a^{-2} \mathcal{F}_T(t) (\vec{\nabla} h_{ij})^2 \right]$$

Stability

$$\mathcal{F}_T := 2 \left[G_4 - X \left(\ddot{\phi} G_{5,X} + G_{5,\phi} \right) \right] > 0$$

$$\mathcal{G}_T := 2 \left[G_4 - 2X G_{4,X} - X \left(H \dot{\phi} G_{5,X} - G_{5,\phi} \right) \right] > 0$$

■ Curvature perturbation

$$\mathcal{L} = a^3 \left[\mathcal{G}_S(t) \dot{\zeta}^2 - a^{-2} \mathcal{F}_S(t) (\vec{\nabla} \zeta)^2 \right]$$

Stability

$$\mathcal{F}_S := \frac{1}{a} \frac{d}{dt} \left(\frac{a \mathcal{G}_T^2}{\Theta} \right) - \mathcal{F}_T > 0, \quad \mathcal{G}_S := \dots > 0$$

$$\text{with } \Theta := -\dot{\phi} X G_{3,X} + 2H G_4 - 8H X G_{4,X} \dots$$

Proof of no-go

TK 1606.05831

- Consider a **non-singular**, spatially flat FLRW solution,

$$a(t) > 0, \quad H, \dot{H}, \dots < \infty \quad (-\infty < t < \infty)$$

- Stability**

$$\mathcal{F}_S = \frac{1}{a} \frac{d}{dt} \underbrace{\left(\frac{a \mathcal{G}_T^2}{\Theta} \right)}_{=: \xi} - \mathcal{F}_T > 0 \quad \Rightarrow \quad \boxed{\frac{d\xi}{dt} > a \mathcal{F}_T > 0} \quad (-\infty < t < \infty)$$

(Scalar stability) (Tensor stability)

$$\xi = \frac{a \mathcal{G}_T^2}{-\dot{\phi} X G_{3,X} + 2H G_4 + \dots} \quad \text{never vanishes}$$

(> 0 , Tensor stability)



in a non-singular universe

($< \infty$, non-singular)

- $\xi(t)$ satisfying these requirements must be ...

Proof of no-go

TK 1606.05831



$$0 < a\mathcal{F}_T < \frac{d\xi}{dt} \rightarrow 0 \quad \text{as } t \rightarrow -\infty \text{ or } t \rightarrow +\infty$$

Non-singular cosmological solutions are stable at any moment in the entire history only if

$$\int_{-\infty}^t a\mathcal{F}_T dt \quad \text{or} \quad \int_t^{\infty} a\mathcal{F}_T dt \quad \text{is convergent}$$

What does $\int_{-\infty} a \mathcal{F}_T dt < \infty$ mean?

- *Disformal* transformation

$$a_E = M_{\text{Pl}}^{-1} \mathcal{F}_T^{1/4} \mathcal{G}_T^{1/4} a, \quad dt_E = M_{\text{Pl}}^{-1} \mathcal{F}_T^{3/4} \mathcal{G}_T^{-1/4} dt$$

- “Einstein frame” — Gravitons propagate along null geodesics

$$\mathcal{L}_E = \frac{M_{\text{Pl}}^2 a_E^3}{8} \left[\dot{h}_{ij}^2 - a_E^{-2} (\vec{\nabla} h_{ij})^2 \right] \quad \begin{array}{l} \text{Creminelli, et al. 1407.8439;} \\ \text{Creminelli, et al. 1610.04207} \end{array}$$

- Past incompleteness in the Einstein frame

$$\int_{-\infty}^t a \mathcal{F}_T dt = \int_{-\infty}^{t_E} a_E dt_E < \infty$$

Affine parameter of a null geodesic

— Geodesically incomplete for the propagation of the gravitons

Multiscalar generalization

- The most general **multiscalar**-tensor theory has not been known

- Generalized multi-Galileon theory: $X^{IJ} := -\frac{1}{2}\partial_\mu\phi^I\partial^\mu\phi^J$

$$\mathcal{L} = G_2(\phi^I, X^{IJ}) - G_{3K}(\phi^I, X^{IJ})\Box\phi^K + G_4(\phi^I, X^{IJ})R + \dots$$

Padilla & Sivanesan 1210.4026

... is *not* the most general one

TK, Tanahashi, Yamaguchi 1308.4798

- 3 new Lagrangians for multiscalar-tensor theory Allys 1612.01972

$$\mathcal{L}_1^{\text{ext}} = A_{[IJ][KL]M}\delta_{\nu_1\nu_2\nu_3}^{\mu_1\mu_2\mu_3}\partial_{\mu_1}\phi^I\partial_{\mu_2}\phi^J\partial^{\nu_1}\phi^K\partial^{\nu_2}\phi^L\partial_{\mu_3}\partial^{\nu_3}\phi^M, \quad \mathcal{L}_2^{\text{ext}} = \dots, \quad \mathcal{L}_3^{\text{ext}} = \dots$$

— most general to date

Multiscalar generalization

Akama, TK 1701.02926

- No-go theorem can be extended straightforwardly to generalized multi-Galileon + 3 new Lagrangians

$$\mathcal{L} = \frac{a^3}{2} \left(\mathcal{K}_{IJ} \dot{Q}^I \dot{Q}^J - \frac{1}{a^2} \mathcal{D}_{IJ} \vec{\nabla} Q^I \cdot \vec{\nabla} Q^J + \dots \right), \quad \phi^I = \bar{\phi}^I(t) + Q^I(t, \vec{x})$$

- Stability: $\mathcal{D} = (\mathcal{D}_{IJ})$ must be positive definite

$$\Leftrightarrow \left(\dot{\phi}^1, \dots, \dot{\phi}^N \right) \mathcal{D} \begin{pmatrix} \dot{\phi}^1 \\ \vdots \\ \dot{\phi}^N \end{pmatrix} > 0 \quad \Leftrightarrow \quad \frac{d\xi}{dt} > a\mathcal{F}_T > 0$$

→ same as in the single-field case

- See Creminelli, et al. 1610.04207 for different proof using EFT

Summary

- All non-singular cosmological solutions are plagued with gradient instabilities at some moment in the entire expansion history in the Horndeski theory — the most general 2nd-order scalar-tensor theory (if graviton geodesics are past and future complete)
 - This can be generalized to theories with **multiple** scalar fields
-

- To evade no-go, one must go *beyond* Horndeski

- A new operator $R^{(3)}\delta N$ (in EFT) that is present in GLPV

Creminelli, *et al.* 1610.04207

Gleyzes, *et al.* 1404.6495

- Higher spatial derivatives, $\zeta\partial^4\zeta$, ...
in Horava gravity/Gao's framework

Pirtskhalava, *et al.* 1410.0882;
TK, Yamaguchi, Yokoyama 1504.05710

Gao 1406.0822