

# Hidden universality in the merger rate of the primordial black hole binaries

Teruaki Suyama

(RESCEU, University of Tokyo)

(A01 連携研究者)

## Refs.


B.Kocsis, TS, T.Tanaka, S.Yokoyama, arXiv:1709.09007 (accepted by ApJ)

# **Plan of my talk**

Brief summary of my achievements in the last fiscal year.


Brief summary of my achievements in this fiscal year.

# Discovery of BH binary by LIGO



# Masses in the Stellar Graveyard


*in Solar Masses*



**LIGO BHs are anomalously heavy?**


# BH spin

$$\chi_{\text{eff}} = \frac{c}{GM} \left( \frac{S_1}{m_1} + \frac{S_2}{m_2} \right) \cdot \frac{L}{|L|} \quad -1 \leq \chi_{\text{eff}} \leq 1$$



**BHs have low spin?**  
**BH spins are misaligned?**

LIGO, NSF, Illustration: A. Simonnet (SSU)



**Clarifying the origin of LIGO BHs has become an important topic in astrophysics and cosmology.**

The seemingly unusual features may suggest that those BHs are primordial.


Editors' Suggestion

PDF

HTML

## Primordial Black Hole Scenario for the Gravitational-Wave Event GW150914

Misao Sasaki, Teruaki Suyama, Takahiro Tanaka, and Shuichiro Yokoyama  
Phys. Rev. Lett. **117**, 061101 (2016) – Published 2 August 2016



A theoretical analysis examines the possibility that the gravitational wave signal (GW150914) detected by LIGO was due to the coalescence of primordial black holes created by the extremely dense matter present in the early Universe.

[Show Abstract +](#)

1603.08338

We proposed a scenario that LIGO BHs are PBHs.

This work was initiated by the discussion done during the mini-workshop held by A01 group (March 10, 2016).

# What are primordial BHs?

PBHs=BHs that formed in the very early Universe

PBHs might comprise all/fraction of dark matter  
(dark matter candidate)

## Popular formation mechanism

Direct gravitational collapse of primordial density  
perturbation

S.Hawking 1971

$$M_{\text{PBH}} \sim \rho H^{-3} \sim 10 M_{\odot} \left( \frac{t}{0.1 \text{ms}} \right)$$

Stellar mass PBHs are formed much before  
BBN era.



# **Two things need to be explained.**

How PBHs formed binaries?

Do their mergers explain the observed merger rate?

# Binary formation in the RD era

THE ASTROPHYSICAL JOURNAL, 487:L139–L142, 1997 October 1  
© 1997. The American Astronomical Society. All rights reserved. Printed in U.S.A.



## GRAVITATIONAL WAVES FROM COALESCING BLACK HOLE MACHO BINARIES

TAKASHI NAKAMURA

Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606, Japan

MISAO SASAKI AND TAKAHIRO TANAKA

Department of Earth and Space Science, Osaka University, Toyonaka 560, Japan

AND

KIP S. THORNE

Theoretical Astrophysics, California Institute of Technology, Pasadena, CA 91125


*Received 1997 April 11; accepted 1997 July 23; published 1997 September 2*



### ABSTRACT


HOs are black holes of mass  $\sim 0.5 M_{\odot}$  they must have been formed in the early uni  
e was  $\sim 1$  GeV. We estimate that in this case in our Galaxy's halo out to  $\sim 50$  kpc there exist  $\sim 5 \times$   
hole binaries the coalescence times of which are comparable to the age of the universe, so that the  
e rate will be  $\sim 5 \times 10^{-2}$  events  $\text{yr}^{-1}$  per galaxy. This suggests that we can expect a few events per  
15 Mpc. The gravitational waves from such coalescing black hole MACHOs can be  
tion of interferometers in the LIGO/VIRGO/TAMA/GEO network. Therefore, the  
HOs can be tested within the next 5 yr by gravitational waves.

*Findings:* black hole physics — dark matter — gravitation — gravitational lensing — C



# Two assumptions (Nakamura et al. 1997)

1. After PBHs are formed (by some mechanism), they distribute uniformly in space (Poisson).



**Initially, PBHs are on the flow of the cosmic expansion.**


2. All PBHs have the same mass

# Binary formation in RD era (Nakamura et al. 1997)

**(The rest is not assumption but physical consequence.)**

As the universe expands, distance between PBHs becomes smaller than the Hubble horizon.


When the PBH energy  $2 M_{BH}$  in the volume  $\sim d^3$  exceeds  $\rho_{rad} d^3$ , the PBHs in pair decouple from cosmic expansion and start to come closer by the gravitational force. (Ioka et al. 1998, Ali-Haimoud et al. 2017)



**This can happen only in the RD era.**

# Binary formation in RD era (Nakamura et al. 1997)

**The nearest third PBH (but distant) gives angular momentum to the bound system.**




Once  $x$  and  $y$  are fixed,  $a$  and  $b$  are determined as

$$a = \frac{1}{f_{\text{PBH}}} \frac{x^4}{\bar{x}^3} \quad b = \frac{x^3}{y^3} a$$

# Binary formation in RD era (Nakamura et al. 1997)

We can compute probability distribution of (a,e).



## Uniform distribution

$$dP = \frac{9}{\bar{x}^6} x^2 y^2 dx dy \quad 0 < x < y < \bar{x}$$


Probability in  $(a, a + da)$  and  $(e, e + de)$

$$dP = \frac{3}{4} f^{3/2} \bar{x}^{-3/2} a^{1/2} e (1 - e^2)^{-3/2} da de.$$

In the paper by Nakamura et al. 1997,  $M_{BH} = 0.5M_{\odot}$  and  $\Omega_{PBH} = \Omega_{DM}$  was considered.

In the paper by Sasaki et al. 2016,  $M_{BH} = 30M_{\odot}$  and the formula was extended to the case  $\Omega_{PBH} < \Omega_{DM}$ .

# Merger event rate (Sasaki et al. 2016)




Consistent with LIGO if  $30M_{\odot}$  PBHs constitute about a few  $\times 0.1\%$  of dark matter.



Recently, the same formula has been used to place upper limit on  $\Omega_{PBH}$  from the LIGO observation.

(Ali-Haimoud, Kovetz, Kamionkowski 2017)



**Stellar mass PBHs as the dominant dark matter is strongly disfavored.**

**Various effects that are ignored have been evaluated in other papers.**

(Ioka et al. 1998, Hayasaki et al. 2009, Sasaki et al. 2016, Eroshenko 2016, Ali-Haimoud et al. 2017)

- Tidal force from outer BHs
- Initial peculiar velocity of PBHs
- Three body collisions
- Additional tidal force from dark matter perturbations
- Encounters of other PBHs (later time effect)
- Tidal force from halos (later time effect)
- Dynamical friction from DM and baryon (later time effect)

**Simple analytical estimation suggest that those effects do not lead to the significant change of the result.**

**We have to keep in mind that these studies adopt the two assumptions.**


To summarize, in 2016, we proposed the PBH scenario for the LIGO events.

... in addition to this achievement...

# 第1回村山杯

# 加速クイズ

I organized "Kasoku" quiz.




# What I did in 2017

# Achievement 1

Classical and Quantum Gravity

ACCEPTED MANUSCRIPT

Primordial black holes – perspectives in gravitational wave astronomy

Misao Sasaki<sup>1</sup> , Teruaki Suyama<sup>2</sup>, Takahiro Tanaka<sup>3</sup> and Shuichiro Yokoyama<sup>4</sup>

Accepted Manuscript online 15 January 2018 • © 2018 IOP Publishing Ltd

[What is an Accepted Manuscript?](#)



Accepted Manuscript PDF

**We wrote a review article about PBHs.**

[+ Article information](#)

[Turn on](#)

[Get permission](#)

[Share this article](#)



[Abstract](#)

Abstract

This is a review article on the primordial black holes (PBHs), with particular focus on the massive ones ( $\geq 10^{15}$  g) which have not evaporated by the present epoch by the

**arXiv: 1801.05235** <sup>22</sup>

# Achievement 2

## How do we test the PBH scenario?

### HIDDEN UNIVERSALITY IN THE MERGER RATE DISTRIBUTION IN THE PRIMORDIAL BLACK HOLE SCENARIO

BENCE KOCSIS<sup>1</sup>, TERUAKI SUYAMA<sup>2</sup>, TAKAHIRO TANAKA<sup>3,4</sup>, AND SHUICHIRO YOKOYAMA<sup>5,6</sup>

<sup>1</sup>Institute of Physics, Eötvös University, Pázmány P. s. 1/A, Budapest, 1117, Hungary;

<sup>2</sup> Research Center for the Early Universe (RESCEU), Graduate School of Science,  
The University of Tokyo, Tokyo 113-0033, Japan

<sup>3</sup> Department of Physics, Kyoto University, Kyoto 606-8502, Japan

<sup>4</sup> Center for Gravitational Physics, Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan

<sup>5</sup> Department of Physics, Rikkyo University, Tokyo 171-8501, Japan and

<sup>6</sup> Kavli IPMU (WPI), UTIAS, The University of Tokyo, Kashiwa, Chiba 277-8583, Japan

*Draft version September 27, 2017*

#### ABSTRACT

Primordial black holes (PBHs) form binaries in the radiation dominated era. Once formed, some fraction of them merge within the age of the Universe by gravitational radiation reaction. We investigate the merger rate of the PBH binaries when the PBHs have a distribution of masses around  $\mathcal{O}(10)M_{\odot}$ , which is a generalization of the previous studies where the PBHs are assumed to have the same mass. After deriving a formula for the merger time probability distribution in the PBH mass plane, we evaluate it under two different approximations. We identify a quantity constructed from the mass-distribution of the merger rate density per unit time and volume  $\mathcal{R}(m_1, m_2)$ ,  $\alpha = -(m_1 + m_2)^2 \partial^2 \ln \mathcal{R} / \partial m_1 \partial m_2$ , which universally satisfies  $0.97 \lesssim \alpha \lesssim 1.05$  for all binary masses independently of the PBH mass function. This result suggests that the measurement of this quantity is useful for testing the PBH scenario.

*Keywords:* gravitational waves – stars: kinematics and dynamics – black hole physics

**Meaning of the hidden university will become clear soon.**

**arXiv:1709.09007**

**B.Kocsis, TS, T.Tanaka, S.Yokoyama**<sup>23</sup>


# How do we test the PBH scenario?

## Main message

**Merger rate distribution in the PBH mass plane has hidden universality and is an interesting observable to address this question.**

**B.Kocsis, TS, T.Tanaka, S.Yokoyama 2017**





**In the future, we will have observed many merger events and will be able to discuss about the distribution in the PBH mass plane ( $m_1, m_2$ ).**

**Can we use the event distribution in the mass plane?**

**In order to investigate what kind of feature appears in the distribution in the mass plane in the PBH scenario, we first generalized the formula to the case of the extended PBH mass function  $f(m_{BH})$ .**

## **Assumption**

In Kocsis et al.2017, we considered the extended mass function which is not so broad ( $\lesssim 10$ ) since it is not clear at all if the same mechanism of the binary formation can still work dominantly for very broad mass function.

Apart from this, we do not assume a specific form of  $f(m)$ .

# Merger event rate distribution in $(m_1, m_2)$ plane

$$\mathcal{R}(m_1, m_2, t) = \frac{1}{2n_{\text{PBH}}} \mathcal{R}_{\text{intr}}(m_1, m_2, t) f(m_1) f(m_2)$$

Observable in the future

Mass function

Probability that given BH pair  $(m_1, m_2)$  form a binary and merge at time  $t$ .

We evaluated  $R_{\text{intr}}$  basically following the approach of Nakamura et al.1997.

## Result

$$\mathcal{R}(m_1, m_2, t) = C \tilde{f}(m_1) \tilde{f}(m_2) (m_1 + m_2)^\alpha$$

$C, \tilde{f}(m)$ : sensitive to the PBH mass function

$$0.97 < \alpha < 1.05$$

# Hidden Universality of $\mathcal{R}(m_1, m_2, t)$

$$\mathcal{R}(m_1, m_2, t) = C \tilde{f}(m_1) \tilde{f}(m_2) (m_1 + m_2)^\alpha$$

$$0.97 < \alpha < 1.05$$

$$\ln \mathcal{R} = \ln C + \ln \tilde{f}(m_1) + \ln \tilde{f}(m_2) + \alpha \ln(m_1 + m_2)$$

$$\begin{array}{c} \downarrow \\ \times \frac{\partial}{\partial m_1} \\ \times \frac{\partial}{\partial m_2} \end{array}$$

**Quantity independent of the mass function!**

# Hidden Universality of $R(m_1, m_2, t)$

## Statement

Construct a quantity  $\alpha$  out of the distribution  $R(m_1, m_2, t)$  as

$$\alpha(m_1, m_2, t) \equiv -(m_1 + m_2)^2 \frac{\partial^2}{\partial m_1 \partial m_2} \ln \mathcal{R}(m_1, m_2, t)$$

Then, the PBH mergers in the present mechanism predict

$$0.97 \lesssim \alpha \lesssim 1.05$$

for any PBH mass function (as long as it is not broad).

**Different formation mechanisms predict different value**

$\alpha \approx 1.43$       PBH binary formation at low redshift.  
(Bird et al. 2016, Clesse, Garcia-Bellido 2016)

$\alpha \sim 4$       Dynamical formation scenario (astrophysics BHs)

# Summary

**GW astronomy has just begun.**

**LIGO might have detected PBHs for the first time.**

**The PBH scenario can be tested in the future by GW data.**