

# **Direct Detection Prospects for the Cosmic Neutrino Background (and other Cosmic Relics)**

**Martin Spinrath**

**National Tsing Hua University, Hsinchu, Taiwan**

**Prospects of Neutrino Physics**

**Kavli IPMU, Tokyo, Japan - 12/04/2019**

**Mostly based on:**

**collaborations with V. Domcke [arXiv:1703.08629] and J. Zurita [arXiv:19???.?????] (our work);**

**And PTOLEMY proposals [arXiv:1307.4738, arXiv:1808.01892],**

**A.J. Long, C. Lunardini, E. Sabancilar [arXiv:1405.7654] (work of other people)**



# Outline

- Introduction
- Resonant Absorption
- Mechanical Forces
- Inverse  $\beta$ -Decay Processes
- Summary and Conclusions



# Outline

- **Introduction**
- **Resonant Absorption**
- **Mechanical Forces**
- **Inverse  $\beta$ -Decay Processes**
- **Summary and Conclusions**




# The Cosmic Neutrino Background

- Produced 1 s after Big Bang (CMB: 379k years)
- Number density:  $330 \text{ cm}^{-3} = 6 n_0$
- Temperature: 1.9 K
- Energy: 0.16 meV
- Velocity:  $10^{-3} - 1 c$
- CNB neutron cross section:  $10^{-27} \text{ pb}$  ( $10^{-63} \text{ cm}^2$ )





# Neutrino Flux Comparison




This talk

[Katz, Spiering 2011]



# The Oldest Picture of the Universe (so far)




[PLANCK, taken from esa.int]



# The Oldest Picture of the Universe in the Future?

[Hannestad & Brandbyge '06]




$m_\nu = (10^{-5} \text{ eV}, 10^{-3} \text{ eV}, 10^{-2} \text{ eV}, 10^{-1} \text{ eV})$  from upper left to lower right


# Didn't we already find it?

**Forbes**    Billionaires    Innovation    Leadership    Money    Consumer    Industry    Lifestyle    F

21,761 views | Feb 28, 2019, 02:00am

## Earliest Signal Ever: Scientists Find Relic Neutrinos From 1 Second After The Big Bang


 **Ethan Siegel** Senior Contributor  
**Starts With A Bang** Contributor Group ①  
[Science](#)  
*The Universe is out there, waiting for you to discover it.*

f 

[<https://www.forbes.com/sites/startswithabang/2019/02/28/earliest-signal-ever-scientists-find-relic-neutrinos-from-1-second-after-the-big-bang/#50b3e913d99c>]



# Didn't we already find it?



**Figure 3:** *Left:* Contours showing  $1\sigma$  and  $2\sigma$  exclusions in the  $\alpha$ - $\beta$  plane for the two redshift bins  $z_1$  and  $z_3$ , both from the BAO data alone and after imposing a CMB prior on  $\alpha$ . *Right:* One-dimensional likelihood of  $\beta$  without (blue) and with (red) the  $\alpha$ -prior for the combined redshift bins. The dashed line is the result after marginalizing over the lensing amplitude  $A_L$ .

[Baumann, Beutler, Flauger, Green, Slosar, Vargas-Magaña, Wallisch, Yèche '18 (Nature Physics '19)]

➡ see also the talk by Masahiro Takada on Monday]




# Didn't we already find it?

- In 2018 >95% confidence from BAO+CMB data
- In fact, in 2015 strong evidence from CMB alone already:


FIG. 3: **Top:** 2D constraints on the jointly varying  $\Lambda\text{CDM}+N_\nu+N_\nu^{\delta\phi}$  parameter space. The constraints on  $N_\nu$  (damping) and  $N_\nu^{\delta\phi}$  (phase shift) are essentially orthogonal. **Bottom:** Constraints from March 2013 *Planck* temperature power spectrum measurements on the number of neutrino species from (1) *blue/solid*: varying  $N_\nu^{\delta\phi}$  while holding  $N_\nu$  fixed at three and (2) *green/dashed*: varying along the physical direction  $N_\nu = N_\nu^{\delta\phi}$ . The constraints assume a Gaussian  $\tau$  prior of mean  $\mu = 0.085$  and width  $\sigma = 0.015$ .

[Follin, Knox, Millea, Pan '15]







# The Other Relics = DM



[M33 rot. curve, Source: Wikipedia]

[Chandra picture of the bullet cluster]




[[https://wiki.cosmos.esa.int/planckpla2015/images/2/2f/A15\\_TT.png](https://wiki.cosmos.esa.int/planckpla2015/images/2/2f/A15_TT.png)]

ct Detection Prospects

Martin Spinrath

11

# Fake(?) WIMP Miracle



**Time to think again?!  
Light WIMPS?**

# Outline

- Introduction
- **Resonant Absorption**
- Mechanical Forces
- Inverse  $\beta$ -Decay Processes
- Summary and Conclusions



# Resonant Absorption

[Weiler '82]

- Similar to GZK cutoff for charged cosmic rays
- Resonant scattering

$$\nu_{\text{UHE}} \bar{\nu}_{\text{CNB}} \rightarrow Z$$

- Dip in energy spectrum expected at  $10^{11}$  GeV
  - Highest energetic neutrinos @IceCube have  $O(10^7)$  GeV
- High energetic Z bursts (not seen so far)



# Outline

- Introduction
- Resonant Absorption
- **Mechanical Forces**
- Inverse  $\beta$ -Decay Processes
- Summary and Conclusions



# The Experiment

[Domcke, MS '17]

- Pendulum in neutrino wind


$$a_\nu \gtrsim \frac{g}{l} d$$

- LIGO-like interferometers

$$a_\nu \gtrsim 10^{-16} \text{ cm/s}^2$$

- Einstein telescope maybe

$$a_\nu \gtrsim 3 \cdot 10^{-18} \text{ cm/s}^2$$



[For more general particle physics applications,  
see Englert, Hild, Spannowsky '17]



# Theory: Scattering

[Domcke, MS '17; see also Duda *et al.* '01, ..., Opher '74]

- Results for the three kinematical cases:

$$a_{G_F^2} = \frac{n_\nu}{2 \bar{n}_\nu} \begin{cases} 3 \cdot 10^{-33} \text{ cm/s}^2 & \text{for (R)} \\ 5 \cdot 10^{-31} (m_\nu / 0.1 \text{ eV}/c^2) \text{ cm/s}^2 & \text{for (NR-NC)} \\ 2 \cdot 10^{-27} (10^{-3} / \beta_{\text{vir}}) \text{ cm/s}^2 & \text{for (NR-C)} \end{cases}$$

- Compare to experimental sensitivity:

$$a_\nu \gtrsim 10^{-16} \text{ cm/s}^2$$

# Other "Winds"

[Domcke, MS '17; see also Duda *et al.* '01]

- Solar neutrinos

$$a_{\text{solar}-\nu} \approx 3 \cdot 10^{-26} \text{ cm/s}^2$$

- Cold WIMP Dark Matter ( $m_{\text{DM}} > 1 \text{ GeV}$ )

$$a_{\text{DM}} \approx 4 \cdot 10^{-30} \left( \frac{(A-Z)^2}{76 A} \right) \left( \frac{\sigma_{X-N}}{10^{-46} \text{ cm}^2} \right) \left( \frac{\rho_{\text{dark(local)}}}{10^{-24} \text{ g/cm}^3} \right) \left( \frac{\beta_X}{10^{-3}} \right)^2 \text{ cm/s}^2$$

- Light WIMP Dark Matter ( $m_{\text{DM}} = 3.3 \text{ keV}$ )

$$a_{\text{light DM}} \approx N_c a_{\text{DM}} \approx 10^9 a_{\text{DM}}$$

[There is also plenty of works on ultralight bosonic DM not based on individual particle scattering, see, e.g., Arvanitaki *et al.* '15; Graham *et al.* '15; Aoki & Soda '16; Pierce *et al.* '18; Morisaki & Suyama '18; Fukuda, Matsumoto & Yanagida '18; ...]



# Improvements and Alternatives

[Domcke, MS '17]

- Sensitivity proportional to  $g$  factor
  - Suspension
  - Space
- Give up on pendulum setup
  - free falling masses and wait
- Alternatives to mechanical force experiment
  - Resonant Absorption
  - Inverse beta decay



# Outline


- Introduction
- Resonant Absorption
- Mechanical Forces
- **Inverse  $\beta$ -Decay Processes**
- Summary and Conclusions




# Inverse Beta Decay

- Lots of Neutrinos around
- Radioactive nuclei, e.g. tritium
- Wait for a neutrino capture
- Goes back to Weinberg

[Weinberg '62]



# Energy Spectrum



[Long, Lunardini, Sabancilar '14]



# Numbers

[Long, Lunardini, Sabancilar '14]

- Number of target nuclei:  $2 \times 10^{25}$  (100 g)
- Rate for Dirac particles (no right-helical neutrinos today):

$$\Gamma_{\text{CNB}}^{\text{D}} = \bar{\sigma} c n_0 N_T \approx 4.06 \text{ yr}^{-1}$$

- Rate for Majorana particles (both helicities equally present):

$$\Gamma_{\text{CNB}}^{\text{M}} = 2 \Gamma_{\text{CNB}}^{\text{D}} \approx 8.12 \text{ yr}^{-1}$$

- Background rate within 0.1 eV of endpoint: 2 Hz



# Current Status (?)

[PTOLEMY '18]

PTOLEMY: A Proposal for Thermal Relic Detection of Massive Neutrinos  
and Directional Detection of MeV Dark Matter

E. Baracchini<sup>3</sup>, M.G. Betti<sup>11</sup>, M. Biasotti<sup>5</sup>, A. Boscá<sup>16</sup>, F. Calle<sup>16</sup>, J. Carabe-Lopez<sup>14</sup>, G. Cavoto<sup>10,11</sup>,  
C. Chang<sup>22,23</sup>, A.G. Cocco<sup>7</sup>, A.P. Colijn<sup>13</sup>, J. Conrad<sup>18</sup>, N. D'Ambrosio<sup>2</sup>, P.F. de Salas<sup>17</sup>,  
M. Faverzani<sup>6</sup>, A. Ferella<sup>18</sup>, E. Ferri<sup>6</sup>, P. Garcia-Abia<sup>14</sup>, G. Garcia Gomez-Tejedor<sup>15</sup>, S. Gariazzo<sup>17</sup>,  
F. Gatti<sup>5</sup>, C. Gentile<sup>25</sup>, A. Giachero<sup>6</sup>, J. Gudmundsson<sup>18</sup>, Y. Hochberg<sup>1</sup>, Y. Kahn<sup>26</sup>, M. Lisanti<sup>26</sup>,  
C. Mancini-Terracciano<sup>10</sup>, G. Mangano<sup>7</sup>, L.E. Marcucci<sup>9</sup>, C. Mariani<sup>11</sup>, J. Martínez<sup>16</sup>, G. Mazzitelli<sup>4</sup>,  
M. Messina<sup>20</sup>, A. Molinero-Vela<sup>14</sup>, E. Monticone<sup>12</sup>, A. Nucciotti<sup>6</sup>, F. Pandolfi<sup>10</sup>, S. Pastor<sup>17</sup>,  
J. Pedrós<sup>16</sup>, C. Pérez de los Heros<sup>19</sup>, O. Pisanti<sup>7,8</sup>, A. Polosa<sup>10,11</sup>, A. Puiu<sup>6</sup>, M. Rajteri<sup>12</sup>,  
R. Santorelli<sup>14</sup>, K. Schaeffner<sup>3</sup>, C.G. Tully<sup>26</sup>, Y. Raites<sup>25</sup>, N. Rossi<sup>10</sup>, F. Zhao<sup>26</sup>, K.M. Zurek<sup>21,22</sup>

Submitted to the LNGS Scientific Committee on March 19<sup>th</sup>, 2018

## Abstract


We propose to achieve the proof-of-principle of the PTOLEMY project to directly detect the Cosmic Neutrino Background (CNB). Each of the technological challenges described in [1, 2] will be targeted and hopefully solved by the use of the latest experimental developments and profiting from the low background environment provided by the LNGS underground site. The




# Another Idea

[MS, Zurita WIP; see also Weiler '01, Mellissinos '99, Müller '87]

- Increase the cross section ( $\sim s$ ) by using a beam
- High energy/intensity muon beams available
- Look for electrons in final state



# CERN M2 Beam Line



[taken from [sba.web.cern.ch/sba/BeamsAndAreas/M2/M2-OperatorCourse.pdf](http://sba.web.cern.ch/sba/BeamsAndAreas/M2/M2-OperatorCourse.pdf)]

# CERN M2 Beam Line

- Beam energy: 150 GeV
- Muon rate:  $1.3 \times 10^7$  /s
- Beam "length": 100 cm
- Treating the beam as fixed target, event rate:

$$R = 1.3 \times 10^9 n_\nu \sigma \frac{\text{cm}}{\text{s}}$$





# Physics Cases (Preliminary)

[MS, Zurita WIP]

Physics Case	Estimated Rate $R$
CNB	$10^{-21}$ /year
Solar $\nu$	$10^{-22}$ /year
Atmospheric $\nu$	$10^{-27}$ /year
Sterile $\nu$ DM	$10^{-28}$ /year
Vanilla WIMP	$10^{-33}$ /year
Resonant WIMP	$10^{-18}$ /year

Other Ideas?





# Why are we so much worse than PTOLEMY?

[MS, Zurita WIP]

- Reminder:

$$\Gamma \sim n_\nu \bar{\sigma} N$$

- CNB number density the same

- Cross sections:

$$\bar{\sigma}_{\text{SZ}} / \bar{\sigma}_{\text{PT}} \sim 10^5$$

- Amount of muons/tritium:

$$N_\mu / N_T \sim 10^{-27}$$



# Outline

- Introduction
- Resonant Absorption
- Mechanical Forces
- Inverse  $\beta$ -Decay Processes
- **Summary and Conclusions**



# Summary and Conclusions

- The CNB is one of the earliest pictures of the universe
- Overwhelming indirect evidence
- But no direct observation so far
- Maybe possible via inverse  $\beta$ -decay (PTOLEMY)
- CNB searches can be DM searches as well
- It is fun to think about other ideas as well...





# Theory: Magnetic Torque

[Domcke, MS '17; see also Duda *et al.* '01, ..., Stodolsky '75]

- Neutrino background splits electron energy levels (spin effect  $\rightarrow$  magnetic effect)

$$a_{G_F}^R = \frac{N_{AV}}{A m_{AV}} \frac{2\sqrt{2}}{\pi} G_F \beta_{\oplus}^{\text{CMB}} \frac{\gamma}{R} \sum_{\alpha=e,\mu,\tau} (n_{\nu_\alpha} - n_{\bar{\nu}_\alpha}) g_A^\alpha$$

- For one flavour

$$a_{G_F}^R \approx 4 \cdot 10^{-29} \frac{n_{\bar{\nu}_\mu} - n_{\nu_\mu}}{2 \bar{n}_\nu} \text{ cm/s}^2$$

- Caveats:
  - Experimentally difficult (magnetic effect)
  - Needs lepton asymmetry



# Theory: Scattering I

[Domcke, MS '17; see also Duda *et al.* '01, ..., Opher '74]

- The basic formula

$$a_{G_F^2} = \Phi_\nu \frac{N_{AV}}{A m_{AV}} N_c \sigma_{\nu-A} \langle \Delta p \rangle$$

- Incoming flux:  $\Phi_\nu$
- #nuclei in 1g test material:  $N_{AV} / (A m_A V)$
- Neutrino-nucleus cross-section:  $\sigma_{\nu-A}$
- Coherence factor:  $N_c$
- Average momentum transfer:  $\langle \Delta p \rangle$





# Theory: Scattering II

[Domcke, MS '17; see also Duda *et al.* '01, ..., Opher '74]

- Neutrinos can come in three kinematics
  - relativistic (R)
  - non-relativistic non-clustered (NR-NC)
  - non-relativistic clustered (NR-C)
- Two important numbers
  - The cross-section:  $\sigma_{\nu-A} \approx 10^{-27} \text{ pb} = 10^{-63} \text{ cm}^2$
  - The coherence factor:  $N_c = \frac{N_{AV}}{A m_{AV}} \rho \lambda_\nu^3 \sim 10^{20}$



# Wind vs. Nudges I

[Domcke, MS '17]

- The scattering rate

$$R = \frac{a G_F^2}{\langle \Delta p \rangle}$$

- Numbers for the CNB

$$R_{(R)} \approx 1 \cdot 10^{-4} \frac{n_\nu}{2 \bar{n}_\nu} g^{-1} s^{-1}$$

$$R_{(NR-NC)} \approx 0.02 \frac{n_\nu}{2 \bar{n}_\nu} \frac{m_\nu}{0.1 \text{ eV}/c^2} g^{-1} s^{-1}$$

$$R_{(NR-C)} \approx 0.4 \frac{n_\nu}{2 \bar{n}_\nu} \frac{0.1 \text{ eV}/c^2}{m_\nu} \left( \frac{10^{-3}}{\beta_{\text{vir}}} \right)^2 g^{-1} s^{-1}$$



# Wind vs. Nudges II

[Domcke, MS '17]

- Solar neutrinos

$$R_{\text{solar}-\nu} \approx 2 \cdot 10^{-9} \text{ g}^{-1} \text{ s}^{-1}$$

- Cold WIMP Dark Matter ( $m_X > 1 \text{ GeV}$ )

$$R_{\text{DM}} \approx 8 \cdot 10^{-3} \left( \frac{100 \text{ GeV}/c^2}{m_X} \right) \left( \frac{\sigma_{X-N}}{10^{-33} \text{ cm}^2} \right) \left( \frac{\rho_{\text{dark(local)}}}{10^{-24} \text{ g/cm}^3} \right) \left( \frac{\beta_X}{10^{-3}} \right) \text{ g}^{-1} \text{ s}^{-1}$$

- Light WIMP Dark Matter ( $m_X = 3.3 \text{ keV}$ )

$$R_{\text{light DM}} \approx 4 \cdot 10^5 \left( \frac{3.3 \text{ keV}/c^2}{m_X} \right)^4 \left( \frac{\sigma_{X-N}}{10^{-42} \text{ cm}^2} \right) \left( \frac{\rho_{\text{dark(local)}}}{10^{-24} \text{ g/cm}^3} \right) \left( \frac{\beta_X}{10^{-3}} \right) \text{ g}^{-1} \text{ s}^{-1}$$



# Helicity Composition

[Long, Lunardini, Sabancilar '14]

- Dirac neutrinos

- left-handed active neutrino:

$$n(\nu_{h_L}) = n_0$$

- right-handed active anti-neutrino:

$$n(\bar{\nu}_{h_R}) = n_0$$

- right-handed sterile neutrino:

$$n(\nu_{h_R}) \approx 0$$

- left-handed sterile anti-neutrino:

$$n(\bar{\nu}_{h_L}) \approx 0$$

- $n_0 = 56 \text{ cm}^{-3}$



# Helicity Composition

[Long, Lunardini, Sabancilar '14]

- Majorana neutrinos

- left-handed active neutrino:

$$n(\nu_{h_L}) = n_0$$

- right-handed active neutrino:

$$n(\nu_{h_R}) = n_0$$

- right-handed sterile neutrino:

$$n(N_{h_R}) \approx 0$$

- left-handed sterile neutrino:

$$n(N_{h_L}) \approx 0$$

- $n_0 = 56 \text{ cm}^{-3}$



# Capture Cross Section

[Long, Lunardini, Sabancilar '14]

$$\sigma_j(s_\nu)v_{\nu_j} = \frac{G_F^2}{2\pi} |V_{ud}|^2 |U_{ej}|^2 F(Z, E_e) \frac{m_p}{m_n} E_e p_e A(s_\nu) (f^2 + 3g^2) ,$$

$$F(Z, E_e) = \frac{2\pi Z\alpha E_e/p_e}{1 - e^{-2\pi Z\alpha E_e/p_e}} ,$$

$$A(s_\nu) \equiv 1 - 2s_\nu v_{\nu_j} = \begin{cases} 1 - v_{\nu_j} & , \quad s_\nu = +1/2 \\ 1 + v_{\nu_j} & , \quad s_\nu = -1/2 \end{cases} \quad \begin{array}{l} \text{right helical} \\ \text{left helical} \end{array}$$

$$\Rightarrow \bar{\sigma} \equiv \frac{\sigma_j(s_\nu)v_{\nu_j}}{A(s_\nu)|U_{ej}|^2 c} \simeq 3.834 \times 10^{-45} \text{ cm}^2 = 3.834 \times 10^{-6} \text{ fb}$$





# Event Rates

[PTOLEMY '13]

- $\beta$ -decay electrons from 100 g tritium:  $10^{16}$  /s
- Fraction within 100 eV of endpoint:  $\sim 2 \times 10^{-7}$
- Fraction within 0.1 eV of endpoint:  $\sim 2 \times 10^{-16}$
- Expected event rate in signal region: 2 Hz
- Expected CNB events:  $O(1)$  /yr

