Dark rate measurement status report

Tokyo University of Science
Tatsushi Kinoshita

26th December, 2019 mPMT meeting


List of the PMTs

	ID	Туре	HV	Negative	Positive
Hamamatsu	HPK	R14374	1250	√	-
HZC new PMT	HZC-B1	XP72B2F	-	-	-
	HZC-B2	XP72B2F	Tuned HV	\checkmark	-
HZC old PMT	HZC-A1	XP82B20	-	-	-
	HZC-A2	XP82B20	Tuned HV	✓ (this week)	-

This week...

• I tuned HV value of HZC-A2 to give almost same gain with HPK, then measured the dark rate.

Dark rate measurement




The temperature dependence of dark rate


- 1st time - HZC-A2 1st time - HPK - HZC-B2 1st time 2nd time 2nd time 2nd time
- In my measurement, the order of rate were HPK ≤ HZC-B2 ≪ HZC-A2, but still more than 200 Hz at 13°C.

Future plan


- To decrease the rate, I will check
 - the time intervals of dark noises
 - the frequency of continuous signal.(for HPK and HZC-B2)




Gain (HPK)




Gain at 13 degree Celsius




Gain at 13 degree Celsius




height [ADC counts]



Gain at 20 degree Celsius




Gain at 20 degree Celsius



HV dependence of gain

 Changing HV supplied to Chinese PMT, I measured 1 p.e. pulse height by illuminating low intensity laser.



Dark rate measurement

- We measured dark rate by F-ADC, keeping the temperature at 13 and 20 °C with incubator.
- We counted the dark noise signals above the threshold 0.3 p.e. and 0.4 p.e.
- To check the consistency, I took the data twice for each PMT.

Setup for dark rate measurement

