

# An Independent Measurement of $H_0$ from Lensed Quasars

*Wednesday, March 25, 2020 11:10 AM (40 minutes)*

Strong gravitational lens systems with time delays between the multiple images are a powerful probe of cosmology, particularly of the Hubble constant ( $H_0$ ) that is key to probing dark energy, neutrino physics, and the spatial curvature of the Universe, as well as discovering new physics. The  $H_0$  Lenses In COSMOGRAIL's Wellspring (H0LiCOW) project has measured  $H_0$  from lensed quasars using deep Hubble Space Telescope and AO imaging, precise time delay measurements from the COSMOGRAIL monitoring project, a measurement of the velocity dispersion of the lens galaxies, and a characterization of the mass distribution along the line of sight. Our latest results from a total of six lenses constrains  $H_0$  to be  $73.3(-1.8,+1.7)$  km/s/Mpc for a flat Lambda CDM cosmology, which is a measurement to 2.4% precision. These results are consistent with independent determinations of  $H_0$  using type Ia supernovae calibrated by the distance ladder method, and are in 3.1-sigma tension with the results of Planck CMB measurements. Combined with the latest distance ladder results from the SH0ES project, we find a 5.3-sigma tension between Planck and late-Universe probes, hinting at possible new physics beyond the standard LCDM model and highlighting the importance of this independent probe.

**Presenter:** WONG, Kenneth (Kavli IPMU)

**Session Classification:** Invited talks