Constraint on the nature of dark matter by high-resolution cosmological simulations and next generation spectroscopy observations

Tomoaki Ishiyama

Chiba University

(Institute of Management and Information Technologies)

18H04337 : JSPS Kakenhi Grant-in-Aid for Scientific Research on Innovative Areas (Why does the Universe accelerate? - Exhaustive study and challenge for the future)

- Constraint on the nature of dark matter by high-resolution cosmological simulations and next generation spectroscopy observations
  - in prep
- Statistical properties of substructures around Milky Way-sized haloes and their implications for the formation of stellar streams
  - Morinaga and Ishiyama et al., MNRAS, 2019

Dwarf spheroidal galaxies (dSph)

- Dark matter dominated
- Old stellar populations
- Little gas
- New (ultra-faint) dSphs have been discovered
- Good candidates to constrain the nature of dark matter by HSC and PFS

Fornax Dwarf Spheroidal

Universal $\Sigma v_{\max} - V_{\max}$ relation

- There is an universal scaling relation between dark halo surface density within a radius $r_{v_{\max}}$ and $V_{\max}$
  - Both halos and “subhalos”
- $\Sigma v_{\max}$ is insensitive to baryonic physics

Hayashi, Ishiyama et al., 2017,
See also Hayashi and Chiba 2015

Aim of our kakenhi project

- Quantifying the distribution and kinematics of dwarf-scale subhalos (10^{7-9} Msun) using high-resolution cosmological N-body simulations
  - Inner profile (surface density)
  - Spatial distribution
  - Kinematics (Line-of-sight, tangential velocities)
- Previous works are limited in a few MW halos and poor mass resolution ($> 10^5$ Msun)
  - Improve to a few tens halos and $\sim 10^{3-4}$ Msun resolution
-> Ultra faint dwarf scale
- Constructing big theoretical templates to compare with future observations
  - Not only CDM, but WDM and SIDM

Simulation suite

All simulations are terminated at $z=0$
halo/subhalo identification: Rockstar (Behroozi+ 2013)

<http://hpc.imit.chiba-u.jp/~ishiytm/db.html>

Name	Model	N	L (Mpc/h)	m_p (Msun/h)	#MW halos
CDM16L	CDM	4096^3	16	5.1×10^3	20~30
CDM8L	CDM	2048^3	8	5.1×10^3	5~10
WDM8L3kev	WDM 3kev	1024^3	8	4.1×10^4	5~10
WDM8L5kev	WDM 5kev	1024^3	8	4.1×10^4	5~10
WDM16L3kev	WDM 3kev	2048^3	16	4.1×10^4	20~30
SIDM8L1S	SIDM*	1024^3	8	4.1×10^4	5~10

SIDM: $\sigma/m=1.0 \text{ cm}^2\text{g}^{-1}$ velocity independent
CDM run with the same initial phase also exists

$\Sigma v_{\max} - V_{\max}$ relation in MW halos

- WDM simulations show lower surface density at $V_{\max} < 20 \text{ km/s}$
- Factor ~ 1.5 difference in $\Sigma v_{\max}$ between CDM and WDM at $V_{\max} \sim 10 \text{ km/s}$
- No significant difference between CDM and SIDM ($\sigma/m=1$, velocity independent)

**How to distinguish
CDM and SIDM ?**

Kinematics

- Can not see significant difference in 3D velocity dispersion
- In the inner part, velocity anisotropy is different?
  - Caution: the number of subhalos in the inner halo is not large
  - See also Nadler+ 2020

Reproducing stellar stream in cosmological context

- Progenitors of streams are accreted at $z=0.5-2.5$

Andromeda stream by HSC (Komiyama+ 2018)

Morinaga and Ishiyama et al., 2019

The impact of filamentary accretion of subhaloes on the shape and orientation of haloes

- Haloes with highly anisotropic accretion become more prolate or triaxial, while haloes with isotropic accretion become more spherical or oblate

Morinaga and Ishiyama, arXiv:1912.11647

Summary and future

- By high resolution cosmological N-body simulations, we are testing whether (how) we can use dSphs to constrain the nature of dark matter
- $\Sigma v_{\max} - V_{\max}$ relation can possibly distinguish CDM and WDM if high precision observations of a number of dSphs are conducted
- To distinguish CDM and SIDM, satellite kinematics plays a key role? But there are large variations in SIDM
- Future:
  - Higher resolution simulations by **Fugaku** supercomputer
  - Larger parameter space
  - Can we use stellar streams to constrain the nature of DM?