

Probing the Genesis of Supermassive Black Holes: Emerging Perspectives from JWST and Expectation toward New Wide-Field Survey Observations

Contribution ID: 4

Type: **Invited**

Little Red Dots and the Rise of Faint, Obscured AGN at $z > 5$

Tuesday 19 November 2024 09:00 (30 minutes)

One of the more surprising results from JWST has been the discovery of faint, broad-line AGN at $z > 5$ with luminosities that are 2-3 dex below those of bright quasars found from the ground. I will discuss recent AGN-related results from the CEERS Survey and what they tell us about the growth of SMBHs in the early universe. This includes the discovery of an actively accreting SMBH at $z=8.67$, which is one of the most distant AGN ever identified. The broad-line AGN identified in our NIRSpec observations are powered by black holes with masses of order $10^7 M_{\text{sol}}$, making them the least-massive BHs known in the early universe. We derive host stellar masses for each AGN, allowing us to place constraints on the BH-galaxy mass relationship in the lowest mass range yet probed in the early universe. Studies in this low-mass regime are key to constraining models of BH seeding and the early growth history of SMBHs. Finally, I will discuss the discovery of a large population of faint, obscured AGN at $z > 5$ known as little red dots (LRDs). Roughly 80% of these sources exhibit broad emission lines in their spectra and our X-ray spectral analysis confirms that they are moderately obscured, with column densities of $\log (n_{\text{H}}/\text{cm}^{-2}) > 23$. The number density of these sources is 2-3 dex above that of bright quasars at $z \sim 5-7$ and 1 dex higher than current samples of X-ray AGN at $z \sim 5$. I will discuss the implications of our findings on the fraction of BH growth that is heavily obscured in the early Universe.

Presenter: KOCEVSKI, Dale (Colby College)