



The Onset of Particle Acceleration from MeV Measurements

Matterhorn

Antimatterhorn




Thomas Siegert

Julius-Maximilians-Universität Würzburg, Germany



17.12.2025, MeV-PeV Frontiers, Tokyo, Japan

The MeV range - A Messenger for Everything



Cosmic-Ray De-Excitation Lines

- Questions: 1) How does particle acceleration work?
 2) What are the abundances in the ISM / SNR / CSM?
 3) What is the low-energy cosmic-ray spectrum?




The Milky Way in MeV Gamma-Rays

Positronium
511 keV +
ortho-Ps

Unresolved
Point
Sources

Astrophys.
Model
Components



INTEGRAL/
SPI
Data Points

Inverse
Compton
Scattering




Nuclear
Decay Lines
 ^{26}Al , ^{60}Fe , ...

The “COMPTEL Bump”

Until ~2013


Search for positron annihilation in flight:

- ➡ Positron sources (injection energy)
- ➡ Dark matter?




“COMPTEL Bump”???

IC (2x e⁻)
IC (optical)
Bremsstrahlung
IC (IR)
IC (CMB)
 π^0 decay



SPI Spatial Fit - Inverse Compton



Background-only
residuals

Bisschoff+2019
(Incl. Voyager data)

$\delta_1 = 0$
(low rigidities)

ISRF(opt) x 10
(central star light)

$z = 8$ kpc
(halo size / diffusion)


$\delta_1 = \delta_2 = 0.5$
(secondary/primary)

GALPROP (v56)
Spatial Models

Testing
systematic
flux variations

SPI Spectral Modelling

GALPROP (v56) Spectral Models



$\delta_1 = \delta_2 = 0.5$:

secondary/primary important in 0.5-8 MeV


ISRF(opt.) $\times 10$:

Too much, too steep

$\delta_1 = 0$:

Too shallow, no turn-over

Comparison to COMPTEL



Power-law: $C_0(E/\text{MeV})^\alpha$

$C_0 = (3.1 \pm 0.3) \times 10^{-6} \text{ ph/cm}^2/\text{s/keV/sr}$

$\alpha = -1.39 \pm 0.09$





Cut-off power-law: $C_0(E/\text{MeV})^\alpha \exp(-E/E_c)$

$\alpha = -0.95 \pm 0.16$

$E_c = 4.9 \pm 1.4 \text{ MeV}$ (Bouchet+2011: $4.0 \pm 1.8 \text{ MeV}$)



Modelling the Galaxy

Inverse Compton Scattering





Modelling the Galaxy

Bremsstrahlung



Electron and ion densities




Modelling the Galaxy

Detected High-Energy Components

About ~25 “good” data points between 50 and 8000 keV


Some gamma-ray lines peak out:
511 keV, ^{60}Fe (1173, 1332 keV), ^{26}Al (1809 keV)



Modelling the Galaxy

Missing Components

Low-Energy Cosmic-Ray De-Excitation Lines




Modelling the MeV Galaxy

Low-Energy Cosmic-Ray De-Excitation Lines

Pros/Cons:

- We have never detected any (only for Solar flares: same mechanism)
- They are at the energy boundaries of typical MeV telescopes (small effective area)
- + There are many of them (look for cumulative effect, not individual lines)
- + They should appear everywhere, plus localised (diffuse ISM, SNRs)



Calculating the LECR Gamma-Rays

Approximations in 2D (projected column densities)

De-excitation line flux
(approx.)
[ph/cm²/s/sr]

Cosmic Ray spectrum

Test with Broken PL & “Carrot” models
adopted from e.g. Indriolo+2009

$$I_{\text{exc}} = N_{\text{H}} \sum_{\text{X}} R_{\text{X}} \int F_{\text{CR}}(E) \sigma_{\text{X}}(E) dE,$$

Cross section

estimated with TALYS
[ver. 2.0; Koning+2023]

Hydrogen column density

$$N_{\text{H}} = 2 \times 10^{22} \text{ cm}^{-2}$$

(Roughly reproducing the
diffuse GeV gamma-rays)

Abundances in ISM

Solar abundance [Lodders 2010]
& Metallicity x2



Tomohiko Oka (JMU) et al.
in prep. (2026)

Then complete 3D modelling!


Calculating the LECR Gamma-Rays

Abundances in the ISM (may vary more than 1 oom!)

$$I_{\text{exc}} = N_{\text{H}} \sum_{\text{X}} R_{\text{X}} \int F_{\text{CR}}(E) \sigma_{\text{X}}(E) dE,$$

A green line points from the R_{X} term in the equation to the text below.

Abundances in ISM
Solar abundance [Lodders 2010]
& Metallicity x2



Tomohiko Oka (JMU) et al.
in prep. (2026)

Then complete 3D modelling!

Calculating the LECR Gamma-Rays

LECR spectrum (the thing we want to know)


$$I_{\text{exc}} = N_{\text{H}} \sum_{\text{X}} R_{\text{X}} \int F_{\text{CR}}(E) \sigma_{\text{X}}(E) dE,$$

Cosmic Ray spectrum

Test with Broken PL & “Carrot” models
adopted from e.g. Indriolo+2009

Broken PL model

Model dependency : Broken PL



Tomohiko Oka (JMU) et al.
in prep. (2026)

Then complete 3D modelling!

$$\phi_p = \begin{cases} N_0 \left(\frac{p(E)}{p(E_1)} \right)^{s_{\text{high}}}, & \text{if } E > E_2 \\ N_0 \left(\frac{p(E_2)}{p(E_1)} \right)^{s_{\text{high}}} \left(\frac{p(E)}{p(E_2)} \right)^{s_{\text{low}}}, & \text{if } E_{\text{cut}} \leq E \leq E_2 \\ 0, & \text{if } E \leq E_{\text{cut}} \end{cases}$$

$$s_{\text{low}} = +1.0 - -3.0$$

Calculating the LECR Gamma-Rays

LECR spectrum (the thing we want to know)


$$I_{\text{exc}} = N_{\text{H}} \sum_{\text{X}} R_{\text{X}} \int F_{\text{CR}}(E) \sigma_{\text{X}}(E) dE,$$

Cosmic Ray spectrum

Test with Broken PL & “Carrot” models adopted from e.g. Indriolo+2009

Carrot model

Model dependency : Carrot



Tomohiko Oka (JMU) et al.
in prep. (2026)

Then complete 3D modelling!

$$\phi'_p = \phi_p + f \times N_0 \left(\frac{p(E_2)}{p(E_1)} \right)^{s_{\text{high}}} \left(\frac{p(E)}{p(E_2)} \right)^{\alpha}$$

$$\alpha = -2.7 - -5.0$$

$$f = 0.01 - 0.1$$

Calculating the LECR Gamma-Rays

Nuclear Excitation Cross Sections (TALYS code)

$$I_{\text{exc}} = N_{\text{H}} \sum_{\text{X}} R_{\text{X}} \int F_{\text{CR}}(E) \sigma_{\text{X}}(E) dE,$$

Cross section
estimated with TALYS
[ver. 2.0; Koning+2023]

For simplicity of illustration:
Use max cross section
at mean energy:



^{12}C : 10 MeV

^{16}O : 20 MeV



Tomohiko Oka (JMU) et al.
in prep. (2026)

Then complete 3D modelling!





What does it help to measure LECR lines?

LECR lines step-by-step: **Step 1: Varying the total power P**

- CR proton spectrum: $\frac{dN}{dE} = P \cdot \frac{\alpha + 2}{E_{\max}^{\alpha+2} - E_{\min}^{\alpha+2}} E^{\alpha}$ Total Power: $\int_{E_{\min}}^{E_{\max}} \frac{dN}{dE} E dE = P$

- Only ^{12}C in the ISM: Strongest line 4.439 MeV

➔ **Get P from one measured line.**





What does it help to measure LECR lines?

LECR lines step-by-step: **Step 2: Varying the index α**

- CR proton spectrum: $\frac{dN}{dE} = P \cdot \frac{\alpha + 2}{E_{\max}^{\alpha+2} - E_{\min}^{\alpha+2}} E^{\alpha}$ Total Power: $\int_{E_{\min}}^{E_{\max}} \frac{dN}{dE} E dE = P$

- Only ^{12}C in the ISM: Strongest line 4.439 MeV



➔ But P is degenerate with α .



What does it help to measure LECR lines?

LECR lines step-by-step: Step 3: Adding other elements

- CR proton spectrum: $\frac{dN}{dE} = P \cdot \frac{\alpha + 2}{E_{\max}^{\alpha+2} - E_{\min}^{\alpha+2}} E^{\alpha}$ Total Power: $\int_{E_{\min}}^{E_{\max}} \frac{dN}{dE} E dE = P$
 - ^{12}C & ^{16}O in the ISM: Strongest lines 4.439, 6.129 MeV
- ➔ Get α/P from only two(!) measured lines!




What does it help to measure LECR lines?

LECR lines step-by-step: **Step 5: Add more lines / elements**

- CR proton spectrum: $\frac{dN}{dE} = P \cdot \frac{\alpha + 2}{E_{\max}^{\alpha+2} - E_{\min}^{\alpha+2}} E^{\alpha}$ Total Power: $\int_{E_{\min}}^{E_{\max}} \frac{dN}{dE} E dE = P$

- All elements in the ISM: many many lines

➔ **Get α and(!) P from three+ measured lines! (+ LECR shape!)**





What does it help to measure LECR lines?

LECR lines step-by-step: Step 5: Add more lines / elements

- CR proton spectrum: $\frac{dN}{dE} = P \cdot \frac{\alpha + 2}{E_{\max}^{\alpha+2} - E_{\min}^{\alpha+2}} E^{\alpha}$ Total Power: $\int_{E_{\min}}^{E_{\max}} \frac{dN}{dE} E dE = P$

- All elements in the ISM: many many lines



➔ Get α and(!) P from three+ measured lines! (+ LECR shape!)



What does it help to measure LECR lines?


LECR lines step-by-step: **Step 6: Carefully check TALYS**

- TALYS does not check if line is forbidden
- ^{16}O L=1-0 is an E0 transition (no gamma-rays!)
- ^{16}O L=1-0 is **internal pair creation (IPC: e^+/e^- pair created!)**
- $^{16}\text{O}^*(6049)$ is not a real γ -ray, but leads to 2.5 MeV positrons




What does it help to measure LECR lines?

Line flux ratios for strongest lines



Supernovae: Timescales

Explosion




Day -20-50

$<10^7$ s

Non-gamma-ray observations


Lightcurve



Day <5000

$<10^9$ s


Remnant



Year <1000

$<10^{11}$ s




Superbubble




Year <10⁷

$<10^{15}$ s

Gamma-ray observations and modelling



Bouchet+2015, ²⁶Al




SN Remnants vs. Galactic Diffuse


Pathway to solve the onset of particle acceleration

Injection at SNRs

$$E_{\text{inj}} \sim E_{\text{break,SNR}}$$





SNRs
= hotspots




Propagation in ISM


$$E_{\text{inj,prop}} \sim E_{\text{break,ISM}}$$



“Inference”





“Measurements”




Internal Pair Creation: Link to 511 keV


$$\text{CR} + \text{X} \rightarrow {}^{16}\text{O}^* \rightarrow e^+/e^-(5.027/2 \text{ MeV})$$



$$\text{CR} + \text{X} \rightarrow {}^{16}\text{O}^* \rightarrow \text{anything}$$






forbidden




Injection Problem

- Use positrons as tracers for acceleration
- Similar idea, plus: Annihilation tells us about efficiency
- Expect an annihilation signal for particles that don't escape






The Milky Way from Radio to Gamma







The Positron Puzzle





What do we see?

Where do the
positrons come from?



Why does it
look like that?













Positron Sources in Space


Positron production is ubiquitous!

- **Massive stars / Novae / SNe Ia&II**
→ Radioactivity from β^+ -decay
- **XRBs / Microquasars**
→ “Compact γ -ray source”; Jets; ...
- **Sgr A***
→ Past AGN activity; Accretion disk
- **Cosmic rays**
→ p-p collisions: Secondary positrons
- **Pulsars**
→ Magnetic field interactions
- **Dark Matter**
→ Decay; Annihilation; Excitation
- **Stars**
→ Stellar flares; e^+ -capture
- **Your ad here**




SNe II






Positron Production Mechanisms




**High-energy
processes!**


Dark matter ...




... annihilation




... decay



... deexcitation



$\gamma\gamma$ pair
production



γ /EM-field
pair production

SNRs: Injection, Escape, and Annihilation

Time-dependent escape of positrons from SNRs
as tracers for acceleration



By measuring the annihilation gamma-rays,
we find the escaping spectrum

$$\underbrace{\frac{\partial f}{\partial t}}_{\text{Phase space density}} + \underbrace{\mathbf{u}(\mathbf{r}, t) \cdot \nabla f}_{\text{Advection term}} - \underbrace{\frac{1}{3} (\nabla \cdot \mathbf{u}) p \frac{\partial f}{\partial p}}_{\text{Adiabatic losses: compression/expansion}} = \underbrace{\nabla \cdot [D(\mathbf{r}, p, t) \nabla f]}_{\text{Spatial diffusion}} + \underbrace{\frac{1}{p^2} \frac{\partial}{\partial p} \left[p^2 D_{pp}(\mathbf{r}, p, t) \frac{\partial f}{\partial p} \right]}_{\text{Reacceleration term}} + \underbrace{\frac{1}{p^2} \frac{\partial}{\partial p} \left[p^2 \left(\frac{dp}{dt} \right)_{\text{loss}} f \right]}_{\text{Energy losses}} + \underbrace{Q(\mathbf{r}, p, t)}_{\text{Source term: } Q = n(\mathbf{r}, t) \lambda F(p)} - \underbrace{\Gamma_{\text{ann}}(\mathbf{r}, p, t) f}_{\text{Annihilation term: } \Gamma = n_X(\mathbf{r}, t) \sigma_{\text{ann}} v}$$

SNRs: Injection, Escape, and Annihilation

Time-dependent escape of positrons from SNRs
as tracers for acceleration




By measuring the annihilation gamma-rays,
we find the escaping spectrum

Link to
nucleosynthesis



Radioactive
decay (β^+)



Source term:

$$Q(\mathbf{r}, p, t)$$




Unique
annihilation
spectrum

Link to
LECR spectrum



Internal pair
creation ($^{16}\text{O}^*$)



Conclusion / Outlook

Measuring the LECR spectrum in SNRs and ISM will:

- + shed light on injection problem
- + estimate the ionisation rate in the Galaxy
- + have links to the 511 keV positron puzzle
- +/- be a major target for COSI and successors
- require high sensitivity and spectral resolution





MeV Gamma-Ray Measurements


Coded-Mask Spectrometer Telescope SPI

INTEGRAL

INTERNational
Gamma-Ray
Astrophysics
Laboratory



Patterns from the
sky



SPI specs:

Energy range:

0.02-8 MeV

FWHM:

2.1 keV @ 511 keV

Angular res.:

2.7° resolution

Line sensitivity ($3\sigma/1\text{Ms}$):

$3 \cdot 10^{-5} \text{ ph cm}^{-2} \text{ s}^{-1} @ 1.8 \text{ MeV}$



Conti. sensitivity ($3\sigma/1\text{Ms}$):

$2 \cdot 10^{-10} \text{ erg cm}^{-2} \text{ s}^{-1} @ 1 \text{ MeV}$

Patterns from the
background


Massive Star Nucleosynthesis

Milky Way in ^{26}Al from massive star ejecta (1.809 MeV)



- 1.2-2.4 M_{Sun} of ^{26}Al in the Galaxy
 - Star formation rate: 4-8 $M_{\odot} \text{ yr}^{-1}$
 - $0.3 \times 10^{43} \text{ e}^+ \text{ s}^{-1}$ along Galactic plane
- Constrain star formation and nucleosynthesis at the same time



Siegert+2023



Massive Stars: 20 years of SPI

How does nucleosynthesis feedback from massive star groups affect the chemical evolution of the Milky Way and the Solar System?

Diehl+2006/2010
Wang+2007/2020
Martin+2009/2012
Kretschmer+2013
Krause+2015/2018
Siegert+2017/2024
Pleintinger+2019/2024



⇒ Milky Way

- $M(^{26}\text{Al}) \approx 2 M_{\odot}$
- $\text{SFR} \gtrsim 4 M_{\odot} \text{ yr}^{-1}$
- $\text{CFR} \gtrsim 0.29 \times 10^{-2} \text{ yr}^{-1}$
- $\text{SNR} = (1.4 \pm 1.1) \times 10^{-2} \text{ yr}^{-1}$

⇒ Single Stars and Star Groups

- Reduced explodability for massive stars
- Clustered star formation
- Frequent superbubble merging


⇒ Solar Vicinity

- Large ^{26}Al flux
- Small $^{60}\text{Fe}/^{26}\text{Al}$ flux ratio
- High latitude emission
- Prominent nearby emission


Star Formation

Galaxy
Milky Way

Feedback



Single Star
 γ^2 -Velorum




Star Group
Perseus OB




Nucleosynthesis




Life of a Positron in Space



Source distribution






Positron Annihilation: $e^+ + e^- \geq 2\gamma$

- Annihilation in Flight:

→ Direct annihilation with $E_{\text{kin}}(e^\pm) \geq 0$




→ $E_{\text{kin}}(e^+) = E_{\text{kin}}(e^-) \approx 0$: 511 keV line







INTEGRAL/SPI 511 keV Portrayal

Disk Annihilation budget: $3.1 \times 10^{43} \text{ e}^+ \text{ s}^{-1}$



**Galactic
centre:
 $0.1 \times 10^{43} \text{ e}^+ \text{ s}^{-1}$**




**Bulge:
 $1.7 \times 10^{43} \text{ e}^+ \text{ s}^{-1}$**

Total Milky Way: $(4.9 \pm 1.7) \times 10^{43} \text{ e}^+ \text{ s}^{-1}$


Compact Objects: Microquasars

Microquasars are Galactic positron sources!



$$\gamma + \gamma \rightarrow e^+ + e^-$$



- e^+ annihilated:
 $\approx 10^{42} \text{ s}^{-1}$
- Duty cycle:
 $\approx 10^{-3}$
- Escape fraction:
 $\approx 20\%$




$10^3 - 10^8$ μ Qs expected in the Milky Way:
Understanding the “inner engine” of accreting black holes!

Compact Objects: Microquasars

V404 Cygni flaring




Siebert+2016b, Nature




Pair-plasma
annihilation light
curve in V404 Cygni

Radio emission associated with pair plasma?



Budgeting Galactic Positrons

Microquasars
52%



Massive Stars
6%

Core-Collapse
Supernovae
4%

Dark matter
<1%?


Pulsars
<1%?

Type Ia Supernovae
37%


In-situ annihilation
Escape from Galaxy
Non-steady state




Upper
contribution
limit



Lower
contribution
limit



More
sources
required







Dark Matter in the Light of 511 keV

Decay, Annihilation, De-excitation

$$F_{511} \propto (\rho_{DM})^n ?$$


- DM density very similar to 511 keV profile
- Predict coherently primary (prompt), secondary (Inverse Compton, synchrotron), and tertiary (positron annihilation) emission for galaxies







From Remnants to Superbubbles

Evolution of Supernova Remnants






Hydro-Simulations of Superbubble Phase






What do Line Shapes tell us?



Guessoum+2005



Cold H₂

Cold H

Warm
neutral

Warm
ionised











Hot
plasma

Line shapes trace
warm and partly
ionised gas:

- $T \approx 7000-40000$ K
- $x_{ion} \approx 2-25\%$



Sources and Sinks



Siebert+2022a

